

CORRECTION

Open Access



Correction to: *PIK3CA* mutation correlates with mTOR pathway expression but not clinical and pathological features in Fibro-adipose vascular anomaly (FAVA)

Yumiko Hori^{1,2}, Katsutoshi Hirose³, Michio Ozeki⁴, Kenji Hata⁵, Daisuke Motooka⁶, Shinichiro Tahara¹, Takahiro Matsui¹, Masaharu Kohara¹, Hiroki Higashihara⁷, Yusuke Ono⁷, Kaishu Tanaka⁷, Satoru Toyosawa³ and Eiichi Morii^{1*}

Correction to: *Diagn Pathol* 17, 19 (2022)
<https://doi.org/10.1186/s13000-022-01199-3>

Following publication of the original article [1], the authors noticed the incorrect article title in the published article. The correct article title is “*PIK3CA* mutation correlates with mTOR pathway expression but not clinical and pathological features in Fibro-adipose vascular anomaly (FAVA)”. The original article has been updated.

Fibroipose vascular anomaly (FAVA). *Diagn Pathol.* 2022;17:19. <https://doi.org/10.1186/s13000-022-01199-3>.

Author details

¹Department of Pathology, Osaka University Graduate School of Medicine, 2-2 Yamadaoka, 565-0871 Suita, Osaka, Japan. ²Department of Central Laboratory and Surgical Pathology, National Hospital Organization, Osaka National Hospital, 2-1-14 Hoenzaka, Chuo-ku, Osaka-shi, Osaka 540-0006, Japan. ³Department of Oral Pathology, Osaka University Graduate School of Dentistry, 1-8 Yamadaoka, 565-0871 Suita, Osaka, Japan. ⁴Department of Pediatrics, Graduate School of Medicine, Gifu University, 1-1 Yanagido, Gifu 501-1194, Japan. ⁵Department of Molecular and Cellular Biochemistry, Osaka University Graduate School of Dentistry, 1-8 Yamadaoka, 565-0871 Suita, Osaka, Japan. ⁶Genome Information Research Center, Research Institute for Microbial Diseases, Osaka University, 3-1 Yamadaoka, 565-0871 Suita, Osaka, Japan. ⁷Department of Radiology, Osaka University Graduate School of Medicine, 2-2 Yamadaoka, 565-0871 Suita, Osaka, Japan.

Published online: 02 May 2022

Reference

1. Hori Y, Hirose K, Ozeki M, et al. *PIK3CA* mutation correlates with mTOR pathway expression but not clinical and pathological features in

The original article can be found online at <https://doi.org/10.1186/s13000-022-01199-3>.

* Correspondence: morii@molpath.med.osaka-u.ac.jp

¹Department of Pathology, Osaka University Graduate School of Medicine, 2-2 Yamadaoka, 565-0871 Suita, Osaka, Japan

Full list of author information is available at the end of the article



© The Author(s). 2022 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.